

E2-2007-56

S. R. Gevorkyan\*, D. T. Madigozhin, A. V. Tarasov,  
O. O. Voskresenskaya

ELECTROMAGNETIC EFFECTS  
AND SCATTERING LENGTHS EXTRACTION  
FROM EXPERIMENTAL DATA ON  $K \rightarrow 3\pi$  DECAYS

Submitted to «Particles and Nuclei, Letters»

---

\*On leave of absence from Yerevan Physics Institute

Геворкян С. Р. и др.

E2-2007-56

Электромагнитные эффекты и извлечение длин рассеяния  
из экспериментальных данных по распаду  $K \rightarrow 3\pi$

Взаимодействие в конечном состоянии в распадах  $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$  рассмотрено в рамках нерелятивистской квантовой механики. Мы показываем, как учесть важнейшие электромагнитные эффекты при анализе экспериментальных данных, используя выражения для амплитуд, полученные ранее. Мы предлагаем соответствующие выражения для поправок к этим амплитудам, справедливые как выше, так и ниже порога  $M_{\pi^0\pi^0} = 2m_{\pi^\pm}$ . Эти формулы могут быть использованы при извлечении длин  $\pi\pi$ -рассеяния из измеренных  $M_{\pi^0\pi^0}$ -спектров.

Работа выполнена в Лаборатории физики частиц ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2007

Gevorkyan S. R. et al.

E2-2007-56

Electromagnetic Effects and Scattering Lengths Extraction  
from Experimental Data on  $K \rightarrow 3\pi$  Decays

The final state interactions in  $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$  decays are considered using the methods of non-relativistic quantum mechanics. We show how to take into account the largest electromagnetic effect in the analysis of experimental data using the amplitudes calculated earlier. We propose the relevant expressions for amplitude corrections valid both above and below the two charged pion production threshold  $M_{\pi^0\pi^0} = 2m_{\pi^\pm}$ , including the average effect for the threshold bin. These formulae can be used in the procedure of pion scattering lengths measurement from  $M_{\pi^0\pi^0}$  spectrum.

The investigation has been performed at the Laboratory of Particle Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2007

The experiments DIRAC and NA48/2 at CERN SPS [1, 2] are able to measure the  $\pi\pi$  scattering lengths difference  $a_0 - a_2$ . Its value is predicted by Chiral Perturbation Theory (ChPT) with a high accuracy [3] ( $a_0 - a_2 = 0.265 \pm 0.004$  in units of inverse pion mass). Thus, the extraction of  $a_0, a_2$  from experimental data with comparable precision becomes an important task. Recently, the NA48/2 experiment has discovered an anomaly (cusp) in the invariant mass  $M_{\pi^0\pi^0}$  spectrum of  $\pi^0$  pairs from  $K^\pm \rightarrow \pi^\pm\pi^0\pi^0$  decays. This anomaly is positioned at the charged pions production threshold  $M_c = 2m_{\pi^\pm}$ , and has been explained by N. Cabibbo [4] as a result of charge exchange reaction  $\pi^+\pi^- \rightarrow \pi^0\pi^0$  in final state. It turns out that investigation of these decays below the charged pions threshold  $M_c$  allows one to extract the scattering lengths with a precision not accessible in other current experiments.

Further development of the model has been done in the works [5–7], where the pions final state interactions were considered with the second order precision in scattering lengths terms. Approach developed in [5] provides the possibility to extract  $a_0, a_2$  from experimental data [2] with a precision compatible to the theoretical prediction one. Nevertheless, there are two problems, requiring a special consideration that can be crucial for scattering lengths extraction from experimental data in the vicinity of the threshold.

The first problem is the accounting for a higher orders in scattering lengths terms and their impact on the value of  $a_0, a_2$ . The estimation made in [5] leads to the corresponding 5% error for  $a_0 - a_2$ . In comparison with the precision of ChPT prediction and with the experimental statistical error this is a rather noticeable uncertainty and its decreasing is very desirable\*.

Another effect that has an impact on  $a_0, a_2$  extraction from experimental data is the electromagnetic interaction of pions in the final state.

This issue is not a trivial task [11, 12], as Coulomb interaction between charged pions\*\* below threshold  $M_c$  leads to formation of bound  $\pi^+\pi^-$  states (pionic atoms  $A_{2\pi^\pm}$ ), and construction of them in the framework of perturbation expansion is a doubtful issue.

---

\*More pessimistic view on impact of higher order terms has given in [7].

\*\*The photons radiation leads only to smearing of Coulomb contribution and will be treated elsewhere.

To solve this problem, we use the technique of non-relativistic quantum mechanics\*. Leaving the strict derivation for elsewhere, we cite here the main result for amplitude of the decay under consideration

$$T = (1 + iv_0 f_0)T_0 + 2iv f_x T_+, \quad f_0 = a_{00}/D, \quad f_x = a_x/D, \quad (1)$$

$$D = (1 - iv_0 a_{00})(1 - 2iva_{+-}) + 2v_0 v a_x^2.$$

Here,  $m_0, m$  are the  $\pi^0, \pi^\pm$  masses, respectively;  $v_0 = \sqrt{M^2 - 4m_0^2}/2m_0$  and  $v = \sqrt{M^2 - 4m^2}/2m$  are the neutral and charged pions velocities;  $T_+$  and  $T_0$  are the «unperturbed» matrix elements of  $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$  and  $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$  decays. The inelastic  $a_x$  and elastic  $a_{00}, a_{+-}$  pion-pion scattering amplitudes in the isotopic symmetry limit are  $a_x = (a_0 - a_2)/3$ ,  $a_{00} = (a_0 + 2a_2)/3$  and  $a_{+-} = (2a_0 + a_2)/6$  [5, 6].

The replacement  $a_i \rightarrow f_i$  has small numerical impact on results of previous calculations done according to [5, 6], but it is crucial for inclusion of the electromagnetic interactions under threshold, where the formation of bound states ( $\pi^+ \pi^-$  atoms) takes place.

The expression (1) includes all successive elastic and inelastic interactions in the  $\pi\pi$  system (to all orders in scattering lengths). Their interaction with a spectator pion can be taken into account at two-loop level [5, 6].

As it was discussed earlier [10], to include the Coulomb interactions in the frameworks of considered approach, it is enough to make the simple replacement in (1):

$$v \rightarrow \tau = iv - \alpha [\log(-2ivmr_0) + 2\gamma + \psi(1 - i\xi)], \quad \xi = \alpha/2v, \quad (2)$$

where  $\gamma = 0.5772$ ,  $\alpha = 1/137$  are Euler and fine structure constants, whereas  $\psi(\xi) = d \log \Gamma(\xi)/d\xi$  is digamma function [13]. The parameter  $r_0$  by its meaning is the strong interaction radius, usually taken as  $r_0 \sim 1/m$ . Later on, we cite it in all expressions, but as we checked, its impact on the results of fitting is small. Strictly speaking, the expression (2) is valid in the region  $vmr_0 \leq 1$ , however for considered kinematics it is a rather good approximation.

To go under threshold one needs to perform the common replacement  $v \rightarrow i\tilde{v}$ :

$$\tau = -\tilde{v} - \alpha \left[ \log(2\tilde{v}mr_0) + 2\gamma + \pi \cot(\pi\tilde{\xi}) + \psi(\tilde{\xi}) \right], \quad (3)$$

$$\tilde{v} = \sqrt{4m^2 - M^2}/2m.$$

---

\*The similar approach to  $K \rightarrow 3\pi$  decays was developed by V. Gribov [8, 9].

Due to asymptotic behavior  $\psi(z) \sim \log(z)$  this expression is finite at threshold  $M_c$ . Moreover, it describes both all bound states (pionium atoms) and electromagnetic interactions leading to unbound states under threshold [10].

Above the threshold the expression (2) acquires the imaginary part. Using the well-known [13] relations for digamma function we obtain

$$\tau = \text{Re } \tau + i\text{Im } \tau,$$

$$\text{Re } \tau = -\alpha \left[ \log(2mr_0) + \gamma + \xi^2 \sum_{n=1}^{\infty} \frac{1}{n(n^2 + \xi^2)} \right], \quad (4)$$

$$\text{Im } \tau = \frac{\pi\alpha}{1 - \exp(-2\pi\xi)}.$$

The above expressions allow one to calculate the electromagnetic corrections in a wide kinematic region. For definiteness let us consider the fitting procedure in NA48/2. In order to extract the values of scattering lengths  $a_0, a_2$  [2], the function

$$F(M^2) = N \int f(a_0, a_2, m^2) \Phi(M^2, m^2) dm^2 \quad (5)$$

was fitted to the experimental mass spectrum. Here,  $f$  is the theoretical distribution from [5], whereas  $\Phi$  describes the experimental set-up resolution and acceptance and is obtained by Monte-Carlo simulation. Integral is replaced with a sum, and decay probability  $f$  is calculated in the middle of every  $M_{\pi^0\pi^0}$  bin with a width  $\delta M^2 = 0.00015 \text{ (GeV}/c)^2$ . The bins are centered in such a way that the threshold  $4m_{\pi\pi}^2$  is placed exactly in the middle of one of them (here we will call this bin the «central» one).

This procedure is precise enough as long as one considers only strong interactions in final state. But the electromagnetic interactions lead to the sharp peak at the threshold, thus for the central bin the averaging of theoretical predictions must be done more carefully.

So, it is convenient to consider the behaviour of the electromagnetic corrections in three separate regions. Up to the lower bound of central bin due to smallness of  $\xi$  in this region from (3) we get

$$\tau = \frac{\pi\alpha}{2} \cot\left(\frac{\pi\alpha}{2\tilde{v}}\right) + \alpha \left[ \log(2\tilde{v}mr_0) + \gamma - \frac{\alpha^2}{4\tilde{v}^2} \varsigma(3) \right], \quad (6)$$

$$\varsigma(3) = \sum_1^{\infty} \frac{1}{n^3} = 1.201.$$

This expression for  $\tau$  has to be used there instead of the velocity formula for charged pions  $\tilde{v} = \sqrt{|M^2 - 4m^2|}/2m$ .

Above the charged pions production threshold beginning from the top bound of central bin the interference between direct  $T_0$  and charge exchange  $T_+$  terms appears (unlike in [4]). As a result, to take into account the electromagnetic effects in this region, one has to add the interference term

$$-4\alpha a_x \left[ \log(2mr_0) + \gamma + \frac{\alpha^2}{4\tilde{v}^2 + \alpha^2} + \frac{1}{2} \log \left( \tilde{v}^2 + (\alpha/3)^2 \right) \right] T_0 T_+ \quad (7)$$

to the square of matrix element.

For these regions the averaging procedure (just a value in the bin's center) exploited in [2] is precise enough.

As to the central bin, where the electromagnetic corrections come from two states of  $\pi^+\pi^-$  pairs (bound or unbound), the more accurate averaging is in use. Confining to the second order terms in scattering lengths, as it is done in previous consideration [5], integrating the square of amplitude (1) in central bin, substituting the relevant velocities (3,4) and dividing result by the square of bin width  $\Delta M$ , we obtain the mean value of correction to the square of matrix element in the central bin in the form

$$\bar{T}^2 = \frac{\pi\alpha^3\zeta(3)}{2v_0v_t^2} T_+^2 - 4\alpha a_x [\log(2v_t m r_0) + \gamma - 0.5] T_+ T_0, \quad (8)$$

$$v_t = v(4m^2 + \Delta M^2/2), \quad v_0 = v_0(4m^2).$$

The first term describes all bound states (pionium atoms). As to the second term, it is a result of interference between the direct «unperturbed» production amplitude  $T_0$  and electromagnetic interactions leading to unbound states. Let us note that in (8) we cite only the main electromagnetic corrections in central bin. We estimated the possible contributions from unbound states ( $\sim T_+^2$ ) and interference between direct amplitude  $T_0$  and bound states ( $\sim T_+ T_0$ ) in central bin, which turn out to be small (less than 1% in comparison with main terms in (8)), and so can be safely neglected.

Numerically, the contribution of the first term (pionium atoms) from (8) into the central bin is near 2.5% of «unperturbed» decay probability that coincides with the prediction evaluated from [14]. Simple estimates show that the interference of direct amplitude  $T_0$  with unbound Coulomb part (second term in (8)) is significant and gives approximately the contribution of the same order as the bound states one.

Now it becomes clear why the fit done in [2] with the pionium contribution as a free parameter leads to the Coulomb contribution, that is about two times larger than can be provided by solely bound states.

In conclusion, let us stressed that at present we have the recipe allowing one to introduce the Coulomb effects into the expressions for the square of decay amplitude (taken, for example, from [5]), averaged correctly over every bin, that can be multiplied by phase space factor, convoluted with the experimental acceptance matrix  $\Phi$  and compared with the experimental data.

We are grateful to V. Kekelidze and J. Manjavidze, who draw our attention to the problem and advice during all the work.

### REFERENCES

1. *Adeva B. et al.* // Phys. Lett. B. 2005. V. 619. P. 50 [arXiv:hep-ph/0504044].
2. *Batley J. R. et al.* // Phys. Lett. B. 2006. V. 633. P. 173 [arXiv:hep-ex/0511056].
3. *Colangelo G., Gasser J., Leutwyler H.* // Nucl. Phys. B. 2001. V. 603. P. 125.
4. *Cabibbo N.* // Phys. Rev. Lett. 2004. V. 93. P. 121801 [arXiv:hep-ph/0405001].
5. *Cabibbo N., Isidori G.* // JHEP. 2005. V. 0503. P. 021 [arXiv:hep-ph/0508130].
6. *Colangelo G., Gasser J., Kubis B., Rusetsky A.* // Phys. Lett. B. 2006. V. 638. P. 187 [arXiv:hep-ph/0604084].
7. *Gamiz E., Prades J., Sciemi I.* hep-ph/0602023.
8. *Gribov V. N.* // Nucl. Phys. 1958. V. 5. P. 653.
9. *Gribov N. N.* // ZhETF. 1961. V. 41. P. 1221.
10. *Gevorkyan S., Tarasov A., Voskresenskaya O.* // Phys. Lett. B. 2007. V. 649. P. 159 (in press) [hep-ph/0612129].
11. *Cabibbo N., Isidori G.* In: Curceoni C., Rusetsky A., Widmann E. (Eds.), arXiv:hep-ph/0610201, P. 52.
12. *Gasser J.* Talk at V Kaon Mini Workshop, CERN, December 12, 2006 (<http://indico.cern.ch>).
13. *Abramowitz M., Stegun I.A.* Handbook of Mathematical Functions. Washington: Dover Publications, 1972, 1037 pp.
14. *Silagadze Z. K.* // JETP Lett. 1994. V. 60. P. 689.

Received on April 12, 2007.

Редактор *В. В. Рудниченко*

Подписано в печать 02.08.2007.

Формат 60 × 90/16. Бумага офсетная. Печать офсетная.

Усл. печ. л. 0,5. Уч.-изд. л. 0,72. Тираж 415 экз. Заказ № 55855.

Издательский отдел Объединенного института ядерных исследований  
141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6.

E-mail: [publish@jinr.ru](mailto:publish@jinr.ru)

[www.jinr.ru/publish/](http://www.jinr.ru/publish/)